



# Stabilizing Effect of Magnetic Helicity on Magnetic Cavities in the Intergalactic Medium


Simon Candelaresi, Fabio Del Sordo



# Intergalactic Bubbles




# M87 in radio



(Churazov et al. 2001)


# MS0735.6+7421 cluster



Hubble (visual) + Chandra (X-ray)

(McNamara and Nulsen 2007)


# Kelvin-Helmholtz Instability



(GRAHAMUK/Wikimedia Commons)


- Bubbles should get disrupted.
- What is the reason for their stability?

# Simulations (hydro)



(Brüggen 2003)

# Simulations (jet inflation)




(Sternberg, Soker 2008)

# Magnetic Helicity

Conservation of magnetic helicity:

$$\lim_{\eta \rightarrow 0} \frac{\partial}{\partial t} \int \mathbf{A} \cdot \mathbf{B} \, dV = 0 \quad \eta = \text{magnetic resistivity}$$

Realizability condition:


$$E_m(k) \geq k|H(k)|/2\mu_0$$




Magnetic energy is bound from below by magnetic helicity.

Can magnetic helicity stabilize intergalactic cavities?

# Numerical Experiments

Full resistive magnetohydrodynamics simulations  
with the PencilCode.




$$\frac{\partial \mathbf{A}}{\partial t} = \mathbf{U} \times \mathbf{B} + \eta \nabla^2 \mathbf{A}$$

$$\frac{D\mathbf{U}}{Dt} = -c_S^2 \nabla \left( \frac{\ln T}{\gamma} \ln \rho \right) + \mathbf{J} \times \mathbf{B}/\rho - \mathbf{g} + \mathbf{F}_{\text{visc}}$$

$$\begin{aligned} \frac{\partial \ln T}{\partial t} &= -\mathbf{U} \cdot \nabla \ln T - (\gamma - 1) \nabla \cdot \mathbf{U} \\ &\quad + \frac{1}{\rho c_V T} (\nabla \cdot (K \nabla T) + \eta \mathbf{J}^2 \\ &\quad + 2\rho\nu \mathbf{S} \otimes \mathbf{S} + \zeta \rho (\nabla \cdot \mathbf{U})^2) \end{aligned}$$

$$\frac{D \ln \rho}{Dt} = -\nabla \cdot \mathbf{U}$$

stratified medium  
hot, under-dense bubble



# Numerical Experiments

box size	$L_{xy}, L_z$	24, 96kpc
bubble radius	$r_b$	8kpc
bubble density	$\rho_b$	$2.5 \times 10^{-26} \text{ gcm}^{-3}$
bubble temperature	$T_b$	$4 \times 10^6 \text{ K}$
medium density	$\rho_0$	$10^{-25} \text{ gcm}^{-3}$
medium temperature	$T_0$	$10^6 \text{ K}$
gravitational acceleration	$g$	$3 \times 10^{-7} \text{ cms}^{-2}$
magnetic field strength	$B_0$	$2.5 \times 10^{-6} \text{ G}$
viscosity	$\nu$	$3 \times 10^{27} \text{ cm}^2 \text{s}^{-1}$
magnetic diffusivity	$\eta$	$9 \times 10^{26} \text{ cm}^2 \text{s}^{-1}$
total time	$t_{\text{end}}$	200 – 250Myr

# Initial Condition: Beltrami Field

$$\mathbf{A} = f(r) A_0 \begin{pmatrix} \cos(yk) + \sin(zk) \\ \cos(zk) + \sin(xk) \\ \cos(xk) + \sin(yk) \end{pmatrix}$$

smoothing function:  $f(r) = 1 - (r/r_b)^{n_{\text{smooth}}}$


inside bubble:  $\nabla \times \mathbf{A} \approx k\mathbf{A}$

→  $E_m \propto A_0^2 k^2$


→  $H_m \propto A_0^2 k$

→ Fix magnetic energy, vary magnetic helicity.


# Initial Condition: Spheromak




# Thermal Emission




# Temperature Iso-Surfaces



hydro




low helicity





high helicity

# Bubble Coherence



# Bubble Coherence



Helical magnetic fields can stabilize the bubbles.

# Conclusions

- Magnetic helicity as constraint on plasma dynamics.
- Magnetic helicity leads to stability at small magnetic energy.
- Mechanism to stabilize intergalactic bubbles.

(Candelaresi and Del Sordo 2020 ApJ **896** 86)

[simon.candelaresi@gmail.com](mailto:simon.candelaresi@gmail.com)